



## **FIRST AID**




If you hear the warning signal of the breathing monitor, check whether the child is crying or moving.



- Stimulate the infant on the bottom of its foot. Keep calling its name.
- If the infant does not respond by any movement or crying, cry out loud for help. If you are not alone, make the other person call the emergency (112 or 999).



- Put the head of the infant to a neutral position. If it is older than one year, tilt its head backwards.
- Make sure the airway is clear of any obstructions. Never search blindly in the child or infant's mouth.




► Monitor breathing for 10 seconds. Place your ear to the mouth of the infant and watch the movement of its chest.







## **FIRST AID**



If the infant is not breathing, place your mouth over both its nose and mouth and breathe in for about 1 second until the chest moves up. Repeat 5 times.



▶ Press 15 x by 2 fingers the middle of the infant's chest between the nipples. Press at the rate of 2 compressions a second to 1/3 of the chest depth.



- ▶ Breathe in twice again, until the chest moves up.
- Repeat 15 chest compressions and 2 breaths.
- After one minute call the emergency (112 or 999), unless another rescuer has done it already.



- If possible, set the **speakerphone mode on** and place the phone next to the head of the infant.
- While dialing, repeat 15 chest compressions and 2 breaths continually.
- Subsequently, follow the instructions of the emergency line operator.

